Philadelphia Area Number Theory Seminar

Farrell Brumley

Universite Paris 13 Nord and Institute for Advanced Study

Concentration properties of theta lifts

Abstract: The classical conjectures of Ramanujan{Petersson and Sato{Tate on the Fourier coe cients of modular forms, or more generally on the Satake parameters of automorphic representations, are highly sensitive to questions of functoriality. For example, the coe cients of CM modular forms are equidistributed according to a very di erent law from that of non-CM forms, and the rst historical counter examples to the naive generalization of the Ramanujan conjecture were found amongst the theta lifts on the group Sp_4 .

A more recent analogue of these conjectures looks at the L^p norms of automorphic forms (with p=1 corresponding to Ramanujan). Their concentration properties, at points or along certain cycles, are of general interest from both an analytic and arithmetic viewpoint. I will describe in this talk a few new results on the subject, joint with Simon Marshall, which attempt to clarify the structure of the problem: the L^p norms of an automorphic form are closely related to the asymptotic size of certain of its periods which in turn re ect the form's functorial origin. In particular, in work in progress, we show the existence of Maass forms, de ned on hyperbolic manifolds and in the image of the theta correspondence from Sp_4 , which concentrate, in a microlocal sense, along closed geodesics.

Wednesday, April 4, 2018 2:40 { 4:00 PM

Bryn Mawr College, Department of Mathematics Park Science Center **328** Tea and refreshments at 2:20PM in Park 339